

THE MECHANICAL MODES OF A 2-PERIODIC TRIANGULATED SURFACE

S. C. POWER

ABSTRACT. A recent "hidden symmetry" conjecture of B. Gin-ge Chen et al is resolved, concerning the dimension of the mechanical modes of a generic 2-periodic triangulated surface $\mathcal{O}$ in $\mathbb{R}^3$ whose structure graph corresponds to a triangular tiling of $\mathbb{R}^2$. We introduce an indexing of the terms in an associated sparse determinant $p_{\mathcal{O}}(z_1, z_2)$ by means of oriented 3-colourings of the underlying sparse graph, and use this and vertex splitting arguments to show that $p_{\mathcal{O}}(z_1, z_2)$ is palindromic or antipalindromic, up to a shift index. This implies the conjectured dimension 1 phenomenon for these surfaces. As a corollary we obtain the topological stability of the generic modes of a 1-periodic triangulated nanotube.

An infinite 2-periodic triangulated combinatorial surface in $\mathbb{R}^3$, considered as a bar-joint framework, may also be viewed as a 2-periodic origami with edge-hinged rigid triangular faces. Such origami, in the case planar structure graphs, as well as more general nontriangulated kirigami, were examined recently by Chen, Liu, Evans, Paulose, Cohen, Vitelli, and Santangelo [2] from the point of view of the analysis of the spectrum of mechanical modes, or rigid unit modes (RUMs). This can be seen as part of a more general endeavour in theoretical physics to understand modes, elasticity and phonons in critically coordinated lattices. See for example Dove et al [4], Kane and Lubensky [5], Lubensky et al [6], Stenull et al [10] and Wegner [11].

We give here a proof that the RUM spectrum of a generic 2-periodic bar-joint framework $\mathcal{O}$ with planar triangulated structure graph is determined by the zero set of a real-valued continuous function on the 2-torus $\mathbb{T}^2$. In particular this resolves a striking conjecture given in [2] that the dimension of mechanical modes of $\mathcal{O}$ is typically equal to 1 rather than 0. This fact is surprising since in the generic case there are no geometric symmetries beyond the given periodicity, and the RUM dimension of general 2-periodic critically coordinated bar-joint framework is expected to be equal to 0. Indeed the dimension here is the algebraic variety dimension for the intersection of the unit 2-torus $\mathbb{T}^2$ with the zero set $V(p_{\mathcal{O}})$ of a complex polynomial $p_{\mathcal{O}}(z_1, z_2)$ with real coefficients. Also this intersection is equal to the intersection of the sets

$$V(\operatorname{Re}(p_{\mathcal{O}})) \cap \mathbb{T}^2, \quad V(\operatorname{Im}(p_{\mathcal{O}})) \cap \mathbb{T}^2$$

and each of these is part of a 1-dimensional curve.

It is known that the dimension 1 property for the RUM spectrum follows from the conjecture in [2] that the origami polynomials $p_{\mathcal{O}}(z)$ are palindromic. This means that, as a function on the 2-torus, the complex conjugate $\overline{p_{\mathcal{O}}(z_1, z_2)}$ agrees with $p_{\mathcal{O}}(z_1, z_2)$ up to a

2000 *Mathematics Subject Classification.* 05C10, 05E99, 52C25, 82D25

Key words: Periodic surface, mechanical mode, vertex splitting, sparse determinant, oriented coloured graph .

multiplicative multinomial factor $z^\alpha = z_1^{\alpha_1} z_2^{\alpha_2}$. We prove that this is true if $\mathcal{O}$ is properly generic. Moreover we show that in the general generic case, for which $\mathcal{O}$ may have linear subframeworks, that $p_{\mathcal{O}}(z)$ is either palindromic or antipalindromic.

The proof has two parts. The first part is a graph-theoretic reduction argument which shows that $\mathcal{O}$ may be obtained from a *base origami*, with a single joint orbit, by a finite sequence of periodic joint-splitting moves. The second part is a combinatorial algebraic argument showing the preservation of a general *palindrome property* under the periodic joint-splitting moves. This property means that $p_{\mathcal{O}}(z)$ is either palindromic or antipalindromic and we show that the latter case holds if and only if the quotient structure graph $G_{\mathcal{O}}$ has an odd number of loop edges.

The origami polynomial is a sparse determinant, that is, the determinant of a sparse function matrix which in this case is determined by a building block motif for $\mathcal{O}$. To analyse such sparse determinants we introduce an indexing of the terms in the determinant expansion with each term being indexed by an oriented 3-colouring of the underlying sparse graph satisfying a coloured outdegree property. This indexing is used to make explicit the effects of bar substitutions under joint splitting, and we expect this formalism to be useful more generally.

The periodic joint-splitting moves correspond to natural vertex-splitting moves on the quotient graph $G_{\mathcal{O}}$. Such quotient graphs, which in general are multi-graphs with loop edges, are realisable on the topological torus in three dimensions, and joint-splitting derives from natural vertex-splitting moves on $G_{\mathcal{O}}$ which respect the facial structure. The local planarity of the graphs G and $G_{\mathcal{O}}$ is captured by local face graphs and diagrams, and we refer to these diagrams to determine the possibilities for colouring extensions associated with each type of joint-splitting move.

It follows from our arguments that the "hidden symmetry" underlying the dimension 1 phenomenon discovered in [2] can be attributed to two factors, namely, (i) the fact that all entries of the matrix function for the base origami are antipalindromic, and (ii) the democratic way in which a periodic joint-splitting move changes the terms of the determinant expansion which sum to the origami polynomial.

We obtain a similar lower dimensional result for the mechanical mode spectrum of a properly generic triangulated 1-periodic cylindrical surface. This covers the setting of 1-periodic nanotubes $\mathcal{N}$. In particular we see that the univariate nanotube polynomial $p_{\mathcal{N}}(z)$ has the palindrome property and so generic points of the spectrum depend stably on perturbation of the (periodic) geometry of $\mathcal{N}$.

The author is grateful to Bryan Chen for introducing him to the palindrome conjecture.

1. BUILDING BLOCK MOTIFS AND POLYNOMIALS.

A 2-periodic bar-joint framework in $\mathbb{R}^3$ is a bar-joint framework in $\mathbb{R}^3$ with two linearly independent *period vectors* a_1, a_2 , which we may assume to lie in the xy -plane. This means that the set of joints and the set of bars are invariant under the group of translations $\mathcal{T} = \{T_k : k \in \mathbb{Z}^2\}$ where

$$T_k(x, y, z) = (x, y, z) + k_1 a_1 + k_2 a_2.$$

We define a *2-periodic triangulated origami* $\mathcal{O}$ to be a 2-periodic bar-joint framework whose infinite underlying (structure) graph G is simple and is associated with a triangulation of the plane. This is the genus zero case of a 2-periodic triangulated surface and we shall confine attention here to this case. It follows that G has a set of *facial 3-cycles* corresponding to the faces of $\mathcal{O}$ and each edge of G is incident to two faces. For our generic considerations there is in fact no loss of generality in assuming that the xy -plane projection of a generic building block motif for $\mathcal{O}$ gives a planar graph, and so the projection of $\mathcal{O}$ is identifiable with G .

An *origami polynomial* $p_{\mathcal{O}}(z_1, z_2)$ for $\mathcal{O}$ is determined by a choice of a building block motif $\mathcal{M} = (F_v, F_e)$ for the given period vectors, and is equal to the determinant of the $3n \times 3n$ square matrix function $\Phi_{\mathcal{O}}(z_1, z_2)$ defined below, where $n = |F_v|$ and $3n = |F_e|$. The motif notation here follows the formalism given in Owen and Power [8], Power [9] and Badri et-al [1]. Thus the joints (resp. bars) of $\mathcal{O}$ are obtained from the translates of the joints (resp. bars) in F_v (resp. F_e) under the action of $\mathcal{T}$. As a bar-joint framework (G, p) the joints of $\mathcal{O}$ appear in the periodic framework vector $p = (p_{\kappa, k})$, where $1 \leq \kappa \leq n$ ranges over the finite index set for the joints in F_v , and $k \in \mathbb{Z}^2$. The edges of G correspond to the bars of $\mathcal{O}$ and we have $p_{\kappa, k} = T_k p_{\kappa, 0}$ for all $k \in \mathbb{Z}^2$.




FIGURE 1. A planar projection of a building block motif with orthogonal period vectors.

The matrix function $\Phi_{\mathcal{O}}(z_1, z_2)$ may be directly specified in terms of a labelling of its rows by the bars $e_1, \dots, e_m$ in F_e and by a labelling of triples of columns by the joints $v_1, \dots, v_n$ in F_v . The row for a bar e of F_e with endpoints $p_{i,k}, p_{j,l}$, with $k, l \in \mathbb{Z}^2$, has at most 6 nonzero entries and these appear in the 6 columns for the pair v_i, v_j . These columns are labelled $v_{i,x}, v_{i,y}, v_{i,z}, v_{j,x}, v_{j,y}, v_{j,z}$. For $i < j$ the row for e takes the following form, where $p(e)$ is the bar vector $p_{i,k} - p_{j,l}$:

$$e \begin{bmatrix} 0 & \cdots & 0 & \overset{v_i}{p(e)z^{-k}} & 0 & \cdots & 0 & \overset{v_j}{-p(e)z^{-l}} & 0 & \cdots & 0 \end{bmatrix}$$

If $i = j$ then the row takes the form

$$e \begin{bmatrix} 0 & \cdots & 0 & \overset{v}{p(e)(z^{-k} - z^{-l})} & 0 & \cdots & 0 \end{bmatrix}$$

and is only determined up to sign. We say that in the case $i = j$ the bar e is *reflexive*. This matrix function is also known as the *symbol function* of $\mathcal{O}$ ([8]).

The origami polynomial associated with the motif is $p_{\mathcal{O}}(z_1, z_2) = \det \Phi_{\mathcal{O}}(z_1, z_2)$. In fact this polynomial is essentially uniquely determined by the choice of period vectors, rather

than the motif $\mathcal{M}$, the only ambiguity being a monomial multiplier $\pm z^k = \pm z_1^{k_1} z_2^{k_2}$. (See [8].) The RUM spectrum $\Omega(\mathcal{O})$ is defined to be the zero set of $p_{\mathcal{O}}(z_1, z_2)$ on the torus $\mathbb{T}^2$. It may be interpreted, on the one hand, as the set of multi-phases of phase-periodic infinitesimal flexes, or, on the other hand, as the set of reduced wave vectors of the zero energy modes in the spectrum of phonon dispersion curves. A proof of this equivalence is given in Power [9]. We remark that despite the dependence of the set $\Omega(\mathcal{O})$ on a choice of motif, the term *spectrum* is appropriate in a technical sense since $\Omega(\mathcal{O})$ may be viewed as the union of the Bohr spectra of almost periodic infinitesimal flexes [1].

From the point of view of a periodic bar-joint framework a periodic origami $\mathcal{O}$ is said to be *generic*, or, more precisely, generic modulo periodicity, if, up to some translation, the coordinates of the motif joints form an algebraically independent set. Thus, paradoxically perhaps, a 2-periodic origami whose motif has a single joint and, necessarily, 3 coplanar incident bars, is generic in this sense, despite being planar. Also, more generally, if $\mathcal{M}$ contains a reflexive bar, then $\mathcal{O}$ contains a set of parallel linear subframeworks. We say that an origami $\mathcal{O}$ is *properly generic* if it is generic and there are no reflexive bars in the motif for $\mathcal{O}$.

Example 1.1. Let $\mathcal{O}$ be a periodic origami, as above, whose motif has a joint set F_v consisting of a single joint. Then $\mathcal{O}$ is referred to as a *base origami*. Suppose that the period vectors are orthogonal unit vectors. Then we may assume that the joint is positioned at the origin, and the bars of the motif have their other joints at $(1, 0)$, $(0, 1)$ and $(1, 1)$. The matrix function for $\mathcal{O}$ is the function $\Phi(z)$ on the torus in $\mathbb{C}^2$ given by

$$\Phi(z_1, z_2) = \begin{matrix} & \begin{matrix} v_{1,x} & v_{1,y} & v_{1,z} \end{matrix} \\ \begin{matrix} e_1 \\ e_2 \\ e_3 \end{matrix} & \begin{bmatrix} 1 - \overline{z_1} & 0 & 0 \\ 0 & 1 - \overline{z_2} & 0 \\ 1 - \overline{z_1 z_2} & 1 - \overline{z_1 z_2} & 0 \end{bmatrix} \end{matrix}$$

It follows that the origami polynomial vanishes identically and the RUM spectrum consists of the entire torus $\mathbb{T}^2$. This is true also for general base origami since these have similar matrix functions.

It is convenient, for diagrammatic depictions, to assume that, as in Figure 1, the geometry is such that $\mathcal{M}$ has a projection which is a planar graph. Let us say that a joint of a bar in F_e which is not in F_v is a *ghost joint*. It is similarly convenient to assume that the motif is chosen so that the ghost joints have one of the three forms $p_{i,(1,0)}$, $p_{i,(0,1)}$ or $p_{i,(1,1)}$.

1.1. Topological stability and palindrome properties. In the context of phonon modes and surface phonons in condensed matter the suggestive term *topologically protected mode* has been used to indicate that upon small periodic perturbation of the geometric parameters the periodic perturbed structure possess a nearby mode with nearby spectral point (wave vector). This is in contrast to the modes that may be present due to internal symmetries and which therefore vanish upon almost all small perturbations. In the analysis of topological insulators for example there is extensive literature concerned with identifying such topological protectedness and the underlying reasons for it.

Similarly, it is of interest to determine the perturbationally stable mechanical modes in a critically coordinated periodic bar-joint structure. One guarantee of such stability, in the case of periodic bar-joint frameworks, is if it happens to be known that in a neighbourhood

of the spectral point λ for the mode the spectrum is a (nonextremal) level curve of a real-valued function. In this way the real-valued determination of the RUM spectrum underlies the existence and persistence of modes under periodic perturbation.

We note that it is elementary to see that a palindromic polynomial $p(z_1, z_2)$ with real coefficients has a zero set on $\mathbb{T}^2$ which is locally determined as the zero set of a real-valued continuous function. Indeed, $p(\overline{z_1}, \overline{z_2}) = z_1^{k_1} z_2^{k_2} p(z_1, z_2)$ for some integers k_1, k_2 , by assumption, and so the function $q(z_1, z_2) = z_1^{k_1/2} z_2^{k_2/2} p(z_1, z_2)$ is real-valued on $\mathbb{T}^2$. A similar argument holds in the antipalindromic case.

2. THE MAIN THEOREM

We now state the main theorem, define various terms, and outline the structure of the proof.

Theorem 2.1. *Let $\mathcal{O}$ be a generic 2-periodic triangulated origami. Then $\mathcal{O}$ has the palindrome property.*

In fact we shall see that $\mathcal{O}$ has a stronger palindrome property associated with subdeterminants of the matrix function $\Phi_{\mathcal{O}}(z)$. Also we show that $p_{\mathcal{O}}(z)$ is palindromic, rather than antipalindromic, if $G_{\mathcal{O}}$ does not have an odd number of loop edges.




FIGURE 2. The face-respecting local action of a periodic joint-splitting move.

A *periodic joint-splitting move*, $\mathcal{O} \rightarrow \mathcal{O}'$, is a construction move which creates a generic 2-periodic triangulated origami $\mathcal{O}'$, with the same period vectors as those of $\mathcal{O}$, such that the joints of $\mathcal{O}'$ include the joints of $\mathcal{O}$, together with a new joint orbit arising from a face-respecting joint-splitting move for the joint v_1 and two incident bars v_1v_2, v_1v_3 . This is indicated, in planar projected form, in Figure 3. In particular, we may define a motif for $\mathcal{O}'$ with $F'_v = F_v \cup \{v_0\}$ and $|F'_e| = |F_e| + 3$. Moreover F'_e is obtained by adding 3 *new bars* incident to v_0 , namely v_0v_1, v_0v_2, v_0v_3 , and by replacing a consecutive set of bars v_1v_j , between v_1v_2 and v_1v_3 , by the *carried bars* v_0v_j .

The proof of the theorem depends on the following two parts, the first of which depends on a construction scheme for *triangulated torus graphs*, as defined below. The second part of the proof requires an analysis of the effect of joint splitting on the origami polynomial.

Part 1. Every generic 2-periodic origami is constructible from a base origami by a sequence of periodic joint-splitting moves.

Part 2. The strong palindromic property, defined in Section 5, is preserved by joint-splitting moves.

We make use of the quotient graph $G_{\mathcal{O}}$ of $\mathcal{O}$ endowed with an inherited facial structure. This is the quotient of the structure graph G of $\mathcal{O}$ with respect to the natural graph automorphisms induced by translation by the periodicity vectors. We may also define it as follows.

The *quotient graph* $G_{\mathcal{M}} = (V, E)$ of $\mathcal{M}$ is the multi-graph with vertex set V equal to F_v , as a set, and where E is obtained from F_e by identifying the joints of bars in F_e with their unique representatives in F_v . Thus $G_{\mathcal{M}}$ is a multi-graph in the sense of having multiple edges and loops. Also $G_{\mathcal{M}}$ inherits a *facial structure* from G in the following sense. The faces of $\mathcal{O}$ are the *edge 3-cycles* $\{e, f, g\}$ given by the unordered triples of bars bordering the triangular faces (or panels) of the bar-joint framework $\mathcal{O}$. The faces of $G_{\mathcal{M}}$ are defined to be the corresponding edge triples. Note however that whereas the faces of $\mathcal{O}$ have distinct triples of joints there may be faces of $G_{\mathcal{M}}$ whose edges have coincident vertices. For examples, the base origami have a quotient graph with a single vertex, 3 edges and 2 faces. The graph $G_{\mathcal{M}}$ is independent of the choice of motif and coincides with $G_{\mathcal{O}}$.

It is evident that $G_{\mathcal{M}}$ is realisable as a graph on the torus with edges represented by smooth curves which do not overlap. To see this one need only construct the torus as the (opposite sides) identification space of the parallelogram defined by the 2-period vectors. In these representations the torus is equal to the closure of the union of connected (open) regions of the torus that are interior to embedded facial 3-cycles of edges.

The quotient graphs so defined are referred to as *triangulated torus graphs*. However, recall that we assume throughout that the structure graph G is a simple graph. It follows that G is 3-connected and it follows from this that $G_{\mathcal{M}}$ is 3-connected.

We thus have the following elementary categorical correspondence.

There is a bijection between 3-connected triangulated torus graphs, defined as identification graphs, and the structure graphs of the periodic triangulated origami $\mathcal{O}$.

In considering the construction of periodic origami by joint splitting moves, it is helpful to consider *face diagrams* which give a planar depiction of the faces of $G_{\mathcal{M}}$ that are incident to v_1 , or a depiction of the faces incident to v_0, v_1 in $\mathcal{G}_{\mathcal{M}'}$, as in Figure 3 for example. Figure 4 indicates that we may also view the joint-splitting move as the result of bar substitutions which derives $\mathcal{M}'$ from $\mathcal{M}$.

If there is no loop edge with vertex v_1 then we may formally define the face graph for vertex v_1 to be the planar graph determined by the faces that are incident to v . If on the other hand v_1 supports one or several loop edges then the incidence of faces to the vertex v_1 can be indicated on an identification space for the torus, or part of the torus.

Figure 5 indicates a vertex splitting move on the base quotient graph which has two faces, one of which is labelled. Figure 6 indicates the results of two vertex splitting moves for a vertex on a loop edge (in the left hand diagram). In the middle diagram the (vertical) loop edge becomes a carried edge (which is not a loop). In the rightmost diagram the loop edge plays the role of one of the split edges and remains in $G_{\mathcal{M}'}$.

It is straightforward to see that in general it is possible for a vertex-splitting move to increase the number of loops by 1.



FIGURE 3. Face graphs for the vertex v_1 in $G_{\mathcal{M}}$, and for the vertices v_1, v_0 in $\mathcal{G}_{\mathcal{M}'}$.

FIGURE 4. Joint splitting as bar substitutions.




FIGURE 5. A torus representation of a vertex-splitting move on the base quotient graph.




FIGURE 6. Two vertex-splitting moves on a loop edge.

3. INDUCTIVE CONSTRUCTION

Lemma 3.1. *Every generic 2-periodic triangulated origami is constructible from a base origami by a sequence of periodic joint-splitting moves.*

Proof. The periodic joint-splitting move provides a natural vertex-splitting move $G_{\mathcal{M}} \rightarrow G_{\mathcal{M}'}$ at the level of quotient graphs. In view of the discussions above it suffices that to

show that if the 3-connected triangulated torus graph $G_{\mathcal{M}'}$ has more than one vertex then it is the result of a vertex-splitting move on a 3-connected triangulated torus graph.

Let v_1, v_2 be distinct adjacent vertices of $G_{\mathcal{M}'}$ and let $F_1 = \{v_1v_2, e, f\}$ and $F_2 = \{v_1v_2, g, h\}$ be the two faces of $G_{\mathcal{M}'}$ that are incident to the edge v_1v_2 . Consider the triangulated torus graph H defined by the operation of edge contraction on $G_{\mathcal{M}'}$ for the edge v_1v_2 . If there are no properly triangulated 3-cycles of edges of the form $C = \{vv_1, v_1v_2, v_2v\}$ then it follows that H is 3-connected.

It remains to consider the case that there does exist an obstacle C to the contraction of v_1v_2 . Such an edge cycle is indicated in Figure 7.




FIGURE 7. A 3-cycle obstruction $C = \{vv_1, v_1v_2, v_2v\}$.

However in this case the graph formed by the triangulation of vv_1v_2 corresponds to a 3-connected triangulation of a sphere. For such triangulations it is well known that there exists an interior edge (one other than v_1v_2, v_1v_5, v_2v_5) which contracts to yield a simple graph which is also a 3-connected triangulation of the sphere or which is a 3-cycle. (See the appendix of [3] for example.) It follows that the contraction of this edge in $G_{\mathcal{M}}$ gives a 3-connected graph K which is the quotient graph of a periodic origami, as required. $\square$

The same argument shows that every properly generic periodic origami is obtained by joint splitting from one which has a motif containing four joints.

Lemma 3.2. *Every properly generic 2-periodic origami is constructible from a properly generic origami with $|F_v| = 4$ by a sequence of periodic joint-splitting moves.*

It is a simple matter to enumerate the four-jointed motifs of the lemma. We remark that, on the other hand, it is a more complicated issue to identify the uncontractible triangulations of the torus which are *strict triangulations* in the sense that there are no 2-cycles or loop edges. For this, see Lawrencenko [7].

4. ORIENTED COLOURINGS AND JOINT SPLITTING

We now introduce a key tool for understanding how the origami polynomial changes under joint-splitting moves, namely the indexing of the terms in the expansion of the determinant of $\Phi_0(z)$ by oriented colourings of the edges of G_0 .

4.1. Oriented colourings and determinants. Recall that a nonreflexive bar e in F_e with endpoints joints $p_{i,k}, p_{j,l}$, for the vertex classes v_i, v_j in F_v and $k, l \in \mathbb{Z}^2$, determines the entry $(p_{i,k} - p_{j,l})_x z^{-k}$ in the column labelled $v_{i,x}$ and the entry $(p_{j,l} - p_{i,k})_x z^{-l}$ in column for $v_{j,x}$. We view the act of choosing one of these two entries as assigning a direction to the bar e , or, equivalently, to its edge in G_M and assigning the label x , or an associated colour to indicate the coordinate choice. Specifically we view the choice of the $v_{i,x}$ entry as assigning the directed edge $v_i v_j$, with v_i as source joint, and the choice of the second $v_{j,x}$ entry as assigning the directed edge $v_j v_i$ with v_j as source.

If, on the other hand, e is a reflexive bar, with $i = j$, with a corresponding loop edge in G_M , then we view the choice of the entry in column $v_{i,x}$ as assigning the label x to e . Also we view e as a directed edge, so that it contributes to the outdegree of v .

Definition 4.1. *An oriented colouring μ of F_e (resp. G_M) is a choice of directions for the bars (resp. edges) and an assignment of labels from the list $\{x, y, z\}$, or from the corresponding colour list $\{\text{blue, red, black}\}$, such that the outdegree of each joint (resp. edge) is 3, with no repeated labels.*

We next relate oriented colourings of F_e to terms in the expansion of the determinant of $\Phi(z) = \Phi_{\mathcal{O}}(z)$.

Note that an oriented colouring μ determines, and is determined by, a choice for each row of $\Phi(z)$ of one of the defined entries of that row in such a way that there are no repeated column choices. Expanding $\det \Phi(z)$ in terms of bijections π between the set of m rows and the set of $m = 3n$ columns, we have

$$\det \Phi(z) = \sum_{\pi: E \rightarrow F_v \times \{x, y, z\}} (-1)^{\text{sgn} \pi} \Phi(z)_{e_1, \pi(e_1)} \cdots \Phi(z)_{e_m, \pi(e_m)}$$

Noting the sparsity of the matrix function $\Phi(z)$ we may omit various zero terms in this sum of $(3n)!$ terms to obtain a sum indexed over the oriented colourings of F_e . Thus

$$\det \Phi(z) = \sum_{\mu \in \mathcal{C}} X_\mu$$

where $\mathcal{C}$ is the set of oriented colourings and where X_μ is the signed $3n$ -fold product of entries of $\Phi(z)$ associated with μ .

If $\mathcal{O}$ is properly generic, meaning that $\mathcal{M}$ has no reflexive bars, then each term X_μ is a nonzero scalar multiple of a monomial z^δ with $\delta \in \mathbb{Z}^2$. (With our assumption on $\mathcal{M}$ the indices δ_1, δ_2 are in fact nonpositive integers.) We define the *monomial degree* of the oriented colouring μ to be $\delta(\mu) = \delta$ in this case. In general however X_μ is a signed scalar multiple of a polynomial, in z_1, z_2 with integral coefficients.

4.2. Local oriented colouring extension. By a *local oriented colouring* λ_1 for a vertex v_1 of G_M we mean a directing and colouring of the edges that have incidence with the vertex v_1 and which is induced by restricting an oriented colouring μ . We write $\lambda_1 \subset \mu$ to indicate this relationship.

In Figure 3 we have indicated local oriented colourings for a vertex-splitting move on a vertex v_1 relative to the incident edges $v_2 v_1, v_3 v_1$. Also we assume that v_1 is not on a loop edge of $\mathcal{G}_M$. Although the joints of F'_v should be labelled differently from those of F_v we find it convenient not to do so in view of the evident correspondence. The left hand part

of Figure 8 indicates a local oriented colouring λ_1 for v_1 , with the three inwardly directed bars to v_1 unlabelled. There are exactly three outwardly directed edges from v_1 with distinct colours, or labels x, y, z , as indicated. Although the μ -inward directed bars in the figure do not have an indicated colour/label their colouring is implied by the μ -colouring of the bars that are not incident to v_1 .

In general there are a number of separate *types* of local oriented colouring depending on the disposition of the three outward bars at v_1 relative to the edges v_1v_2, v_1v_3 of the joint-splitting move. We identify these various types below. We also identify the ways in which an oriented colouring μ may be extended to an oriented colouring μ' of $G_{\mathcal{M}}$ by assigning colours to the new edges and the carried edges. We shall see that in all cases there are 3 edges that may be coloured freely, and so in all cases there are 6 oriented colouring extensions μ' of μ .




FIGURE 8. An oriented colouring extension.

In left hand diagram of Figure 8, for example, which we call type (a), the oriented colouring directions of μ is such that two of these outward edges, namely those for v_1v_2 and v_1v_3 , are edges to be split. The other outward edge is a *fixed edge* representing a fixed bar for the associated joint-splitting move. The right hand diagram in Figure 8 indicates a local colouring extension μ' in which the 3 new edges, v_0v_1, v_0v_2, v_0v_3 may be freely coloured, or labelled a, b, c , to satisfy the outdegree condition, while the carried bars, which are directed inwards, must inherit their direction and labelling.

In Figure 9 we give face graph indications of the four other possible types of local oriented colourings (modulo reflected cases) that are possible, under the assumption, once again, that v_1 is not on a loop edge. In these diagrams we have not indicated further inward edges to v_1 which may be present. We call these types (b), (c), (d) and (e), respectively. Case (b), for example, indicates that μ has the three outward edges from v_1 strictly to one side of the two bars to be split. In case (d) there are two outward edges on one side and none on the other, and so on.

Since we are concerned with generic periodic origami frameworks it should be clear that there are no further types that we need consider (when v_1 does not lie on a loop edge). For example, if the 3 outward directed edges from v_1 are identified as carried edges, so that at the level of the move $\mathcal{O} \rightarrow \mathcal{O}'$ the corresponding bars are carried bars, then the framework $\mathcal{O}'$ is an alternative generic realisation of the framework obtained by a joint-splitting move

of type (b). In particular the origami polynomials share the property of being palindromic, or antipalindromic, or neither of these.




FIGURE 9. Types (b), (c), (d), (e) of outward edge triples.

For types (d), (e), in which there are outward directed edges from v_1 on both side of the split edges, we see that in contrast to the (a), (b) and (c) types, the freely colourable edges are given by 2 new edges and a carried edge (rather than the edge V_0v_1).




FIGURE 10. Type (d) joint splitting and oriented colouring extension.

We now summarise the construction of oriented colouring extensions μ' of μ for all types of μ when v_1 does not lie on a loop edge.

Step (I). The colour and direction of the edges of $G_{\mathcal{M}}$ determine the μ' colour and direction for the following edges in $G_{\mathcal{M}'}$:

- (i) The fixed edges v_1v_j in $G_{\mathcal{M}'}$ (those that are not new edges or carried edges) are μ' -coloured and directed as their corresponding edges in $G_{\mathcal{M}}$.
- (ii) The carried edges v_jv_0 in $G_{\mathcal{M}'}$ are μ' -coloured and directed as their corresponding edges v_jv_1 in $G_{\mathcal{M}}$ if those bars have v_j as a source with respect to μ .
- (iii) If μ is of type (d) or (e) then the edge v_1v_0 in $G_{\mathcal{M}'}$ is μ' -directed with v_1 as source and coloured according to the μ -colour of the remaining carried edge v_1v_k in $\mathcal{M}$.

Step (II). The remaining 3 edges of $G_{\mathcal{M}'}$ (which are v_0v_2, v_0v_3 and either v_0v_1 or v_0v_k) are coloured freely with three distinct colours and directed with v_0 as source.

It follows that the set $\mathcal{C}'$ of all oriented colourings for $G_{\mathcal{M}'}$ is equal to the set of extension colourings μ' . Moreover the latter set is partitioned into disjoint sets according to their associated colouring μ .

In an exactly similar way one can identify the local oriented colouring extensions in the case of joint splitting for a joint on a reflexive bar. The local planar face diagrams now take a different form, with cut faces and identification lines as in Figure 6. Nevertheless the joint-splitting move is a local face-respecting move, with carried bars, fixed bars and 3 new bars, and the colouring extensions are given as before.

Example 4.2. Consider first the vertex-splitting move on the base quotient graph given in the earlier face diagram in Figure 5. In terms of bar substitutions at the level of the motif the joint-splitting move $\mathcal{M} \rightarrow \mathcal{M}'$ corresponds to the replacement of a single reflexive bar by 4 bars which are coincident at the new joint v_0 . There are 6 terms X_μ each of which is zero since one of the reflexive bars is coloured black (labelled z). Thus the origami polynomial is zero, as we have seen. Let μ' be an oriented colouring of μ where the two remaining reflexive bars are coloured blue and red (by μ and μ'). These coloured and directed bars provide together a palindromic factor to $X_{\mu'}$ while the remaining bars provide monomial factors. Thus $X_{\mu'}$ is palindromic. Moreover the shift index is independent of the various extensions μ' and also independent of the two red-blue choices for the two reflexive bars. It follows that $p_{\mathcal{O}'}(z)$ is palindromic.

One can similarly examine a joint-splitting move such as that indicated in Figure 11. Reassigning the $\mathcal{M}$, $\mathcal{M}'$ notation above, this move ($\mathcal{M} \rightarrow \mathcal{M}'$) leaves a single reflexive bar in $\mathcal{M}'$. Once again the move is effected by replacement of a single reflexive bar by 4 nonreflexive bars, as before. The substitutions in this case lead to an antipalindromic origami polynomial.




FIGURE 11. Vertex splitting on $G_{\mathcal{M}}$ leaving a single loop edge in $G_{\mathcal{M}'}$.

5. PART 2 OF THE PROOF

Recall once again that the determinant $\det \Phi_{\mathcal{O}}(z)$ is expanded as the sum of terms X_μ each of which is a scalar multiple of a product of factors of the form z_1^{-1} , z_2^{-1} , $z_1^{-1}z_2^{-1}$, $(1 - z_1^{-1})$, $(1 - z_2^{-1})$ and $(1 - z_1^{-1}z_2^{-1})$. We write $\text{pol}(\mu)$ for this product and note that it is either palindromic or antipalindromic, in the sense of Definition 5.1

In the light of the previous section it is now straightforward to identify the various changes

$$X_\mu \rightarrow X_{\mu'}$$

for a joint splitting move $\mathcal{M} \rightarrow \mathcal{M}'$. The term X_μ is a signed product of $3n$ matrix entries of $\Phi_{\mathcal{O}}(z)$. For an oriented colouring extension μ' of μ we derive $X_{\mu'}$ from X_μ by noting the local bar substitutions associated with the joint-splitting move. In all cases the product

$X_{\mu'}$ is obtained from the product X_{μ} by Steps I and II above, which, more informally, amount to

- (i) replacing the terms for bars that are to be "carried", by the terms of the carried bars (with their possibly new μ' colouring),
- (ii) including 3 new multiplicative factors corresponding to the new bars, and
- (iii) making the appropriate permutation sign change.

Definition 5.1. A polynomial $p(z)$ in the commuting variables $z_1, z_2, z_1^{-1}, z_2^{-1}$ is *palindromic* (resp. *antipalindromic*) if $p(z) = z^{\alpha} \overline{p(z)}$ (resp. $p(z) = -z^{\alpha} \overline{p(z)}$) for some monomial $z^{\alpha}, \alpha \in \mathbb{Z}^2$. The index α is the shift index for $p(z)$.

Note that these notions also extend to rational functions in the variables.

Let λ be a local oriented colouring for the joint v in F_v and let

$$X^{\lambda} = \sum_{\mu: \lambda \subset \mu} X_{\mu}$$

Definition 5.2. Let $\mathcal{O}$ be a generic periodic origami and let v be a joint of $\mathcal{M}$. Then

- (i) $\mathcal{O}$ has the local palindrome property at v if there is a multi-index $\alpha = (\alpha_1, \alpha_2)$ and a choice $c \in \{1, -1\}$ such that $X^{\lambda} = cz^{\alpha} \overline{X^{\lambda}}$ for every local oriented colouring λ for v .
- (ii) $\mathcal{O}$ has the strong palindrome property if $\mathcal{O}$ has the local palindrome property at v for every joint.

Since $p_{\mathcal{O}}(z)$ is the sum of the polynomials X^{λ} over all the local oriented colourings of v it follows that property (i) implies that the origami polynomial is palindromic or antipalindromic.

Lemma 5.3. Let $\mathcal{O}$ be a generic periodic origami and let $\mathcal{O} \rightarrow \mathcal{O}'$ be a periodic joint-splitting move. If $\mathcal{O}$ has the strong palindrome property then so does $\mathcal{O}'$.

Proof. Suppose first that the joint-splitting move is with respect to a joint which does not lie on a reflexive bar.

Fix μ and λ_1 with $\lambda_1 \subset \mu$ a local oriented colouring for the joint v_1 . Let μ' be an oriented colouring extension of μ . There are 6 such extensions $\mu'_i, 1 \leq i \leq 6$, associated with the colour choices for the 3 freely colourable bars indicated in Step (II) of the previous section. A specific colour choice, say the i^{th} , provides an oriented colouring choice, which we shall denote $\lambda'_{1,0,i}$, for the set of bars in $\mathcal{M}'$ that have v_0 or v_1 as an end joint.

Since the bars in $\mathcal{M}$ incident to the joint v_0 are also not reflexive bars it follows that the polynomial factor $\text{pol}(\mu'_i)$ of $X_{\mu'}$ is equal to the product of $\text{pol}(\mu)$ and a fixed monomial factor which is independent of i . Thus

$$X_{\mu'_i} = d_i X_{\mu}$$

where d_i is a scalar multiple of a monomial whose monomial degree $\beta \in \mathbb{Z}^2$ is independent of i . The scalar factor in d_i depends on i according to the colouring of the bars of $\mathcal{M}$ and $\mathcal{M}'$ that feature in the bar substitution associated with the joint-splitting move. Thus we obtain

$$\sum_{\mu': \lambda'_{1,0,i} \subset \mu'} X_{\mu'} = d_i \sum_{\lambda_1 \subset \mu} X_{\mu}$$

We now use these equalities and the strong palindrome property for $\mathcal{O}$ to deduce the strong palindrome property for $\mathcal{O}'$.

Let $\lambda'_{1,j}$ be the local oriented colouring of $\mathcal{M}'$ induced by restricting $\lambda'_{1,0,j}$ to the bars with v_1 as an end joint. For types (a), (b), (c) there are 2 values of i for which $\lambda'_{1,0,i}$ restricts to a $\lambda'_{1,j}$. This is because in each case the bar with joints v_0, v_1 is one of the freely colourable bars. For types (d), (e) on the other hand there are 6 values of i . In the case of a 2 values type we have

$$\begin{aligned} X^{\lambda'_{1,j}} &= \sum_{i=i_1, i_2} \left(\sum_{\mu': \lambda'_{1,0,i} \subset \mu'} X_{\mu'} \right) \\ &= (d_{i_1} + d_{i_2}) \sum_{\mu: \lambda_1 \subset \mu} X_{\mu} \\ &= (d_{i_1} + d_{i_2}) X^{\lambda_1} \end{aligned}$$

Since d_{i_1}, d_{i_2} are multiples of a single monomial, z^β say, it follows that $X^{\lambda'_{1,j}}$ is palindromic with combined shift index $\alpha + \beta$. This argument also holds for μ of type (a), (b) or (c), the only difference being that there are 6 factors $d_1, \dots, d_6$.

To see that the local palindrome property holds at v_0 the argument is similar. Note first that for all types (a), (b), (c), (d), (e), if μ' and μ'' are both extensions of μ , and if μ' and μ'' agree on the bars with v_0 as an end joint, then $\mu' = \mu''$. Let $\lambda'_{0,j}$, $1 \leq j \leq 6$, denote the restrictions of μ' to the bars with v_0 as an end joint. We have

$$\begin{aligned} X^{\lambda'_{0,j}} &= \sum_{\mu': \lambda'_{1,0,j} \subset \mu'} X_{\mu'} \\ &= d_j X^{\lambda_1} \end{aligned}$$

and the local palindrome property at v_0 follows.

For joints of F'_v , other than v_1, v_0 , it similarly follows that the local palindromic property holds.

Suppose now that v_1 lies on a reflexive bar. Then the argument above applies with the following routine changes. The factors d_i are now scalar multiples of a rational function $q_i(z)$, which is independent of i , and we have $\text{pol}(\mu') = q_i(z) \text{pol}(\mu)$. Once again, $q_i(z)$ is either palindromic or antipalindromic, as is $\text{pol}(\mu')$. $\square$

6. 1-PERIODIC TRIANGULATED CYLINDRICAL SURFACES

We consider now an infinite 1-periodic triangulated surface in $\mathbb{R}^3$, considered once more as a bar-joint framework $\mathcal{N} = (H, p_H)$, in the setting where the structure graph H is simple and is associated with a triangulation of the infinite cylinder $\mathbb{R} \times S^1$. We refer these frameworks as triangulated nanotubes, even though their girth may be large. The methodology for the determination of the RUM spectrum in $\mathbb{T}$ closely follows that in Section 1 except that there is now one periodicity vector a_1 with translation group

$$(x, y, z) \rightarrow (x, y, z) + k a_1, \quad k \in \mathbb{Z},$$

and the symbol function is a univariate function matrix, $\Phi_{\mathcal{N}}(z_1)$.

One can obtain palindromicity properties for the generic univariate nanotube polynomial as a corollary to Theorem 2.1. Indeed, note that there is a nongeneric 2-periodic origami $\mathcal{O} = (G, p)$ that can be viewed as being wrapped onto $\mathcal{N}$ while respecting 1-periodicity and bar incidence. Here G is a natural covering graph for H and one may identify the respective quotient graphs $H_{\mathcal{N}}$ and $G_{\mathcal{O}}$. It follows readily that

$$\Phi_{\mathcal{N}}(z_1) = \Phi_{\mathcal{O}}(z_1, 1), \quad p_{\mathcal{N}}(z_1) = p_{\mathcal{O}}(z_1, 1)$$

If $\mathcal{N}$ has an even number of linear subframeworks then $H_{\mathcal{N}}$ has an even number of loop edges (and conversely). Thus $G_{\mathcal{O}}$ has an even number of loop edges and so by the analysis in Section 5 $p_{\mathcal{O}}(z_1, z_2)$ is palindromic. Thus $p_{\mathcal{N}}(z_1)$ is palindromic.

Theorem 6.1. *Let $\mathcal{N}$ be a generic 1-periodic triangulated nanotube and let r be the number of linear subframeworks (or loop edges in the quotient structure graph). Then the univariate nanotube polynomial $p_{\mathcal{N}}(z_1)$ is palindromic if r is even and is antipalindromic if r is odd.*

REFERENCES

- [1] G. Badri, D. Kitson and S. C. Power, The almost periodic rigidity of crystallographic bar-joint frameworks, *Symmetry* 6 (2014), 308-328.
- [2] B. Gin-ge Chen, B. Liu, A. A. Evans, J. Paulose, I. Cohen, V. Vitelli, and C.D. Santangelo, Topological mechanics of origami and kirigami, arXiv:1508.00795v2.
- [3] J. Cruickshank, D. Kitson and S. C. Power, The generic rigidity of triangulated spheres with blocks and holes, *J. Combinatorial Theory B*, to appear.
- [4] M. T. Dove, A. K. A. Pryde, V. Heine and K. D. Hammonds. Exotic distributions of rigid unit modes in the reciprocal spaces of framework aluminosilicates, *J. Phys., Condens. Matter* 19 (2007) doi:10.1088/0953-8984/19/27/275209.
- [5] C. L. Kane and T. C. Lubensky. Topological boundary modes in isostatic lattices. *Nature Phys.*, 10(1):39-45, 2014.
- [6] T. C. Lubensky, C. L. Kane, Xiaoming Mao, A. Souslov, Kai Sun, Phonons and elasticity in critically coordinated lattices, arXiv:1503.01324.
- [7] S. Lawrencenko, Irreducible triangulations of a torus, *Ukrain. Geom. Sb.* 30 (1987), 52-62.
- [8] J. C. Owen and S. C. Power, Infinite bar-joint frameworks, crystals and operator theory, *New York Journal of Mathematics*, 17 (2011), 445-490.
- [9] S. C. Power, Polynomials for crystal frameworks and the rigid unit mode spectrum, *Phil. Trans. R. Soc. Lond. Ser. A Math. Phys. Eng. Sci.* 372 (2014), DOI: 10.1098/rsta.2012.0030.
- [10] O. Stenull, C. L. Kane and T. C. Lubensky, Topological phonons and Weyl lines in 3 dimensions, arXiv:1606.00301.
- [11] F. Wegner, Rigid-unit modes in tetrahedral crystals, *J. Phys.: Condens. Matter* 19 (2007) 406218 doi: 10.1088/0953-8984/19/40/406218.

DEPT. MATH. STATS., LANCASTER UNIVERSITY, LANCASTER LA1 4YF, U.K.
E-mail address: s.power@lancaster.ac.uk